

Small Gamma Products with Simple Values

Albert Nijenhuis

Introduction. Central as the gamma function is, it is surprising that its only known simple specific values are $\Gamma(m)$ and $\Gamma(m + \frac{1}{2})$ for integral m . There are, however, numerous formulas that relate specific values of Γ to other functions, such as elliptic or hypergeometric functions - or to other specific values of Γ itself. Among the latter there are products that have very simple values. An example is

$$(1) \quad \Gamma\left(\frac{1}{14}\right) \Gamma\left(\frac{9}{14}\right) \Gamma\left(\frac{11}{14}\right) = 4\pi^{3/2},$$

which recently occurred in the Problems section of the “Monthly” [1], see also [2]. There is also, of course, the classical multiplier formula

$$(2) \quad \prod_{k=1}^{m-1} \Gamma\left(\frac{k}{m}\right) = (2\pi)^{(m-1)/2} m^{-1/2},$$

but it would be more interesting to find simple values for products of fewer factors. In this note we do just that, for a large class of products, and with little computational effort.

Consider any odd integer $n > 1$ and the set $\Phi(2n)$ of numbers in $[0, 2n]$ that are relatively prime to $2n$. (Its cardinality is $\varphi(2n)$, Euler’s totient function.) $\Phi(2n)$ is a group with respect to multiplication modulo $2n$. Let $\nu(n)$ be the order of the subgroup generated by $n + 2$, and let A be this subgroup or any one of its cosets. Let $b(A)$ count the $x \in A$ that are larger than n . Our main result is

Theorem.

$$(3) \quad \prod_{x \in A} \Gamma\left(\frac{x}{2n}\right) = 2^{b(A)} \pi^{\nu(n)/2}.$$

A few cases of this formula are known, but even for small n there are some gaps in published lists.

The Formula. Throughout this paper, $n > 1$ will denote a “fixed” odd integer, and let $\Phi(n)$ be the set (group) of all integers in the interval $[0, n]$ relatively prime to n . Then $\phi(n) = \phi(2n)$, and the map $\alpha : \Phi(n) \rightarrow \Phi(2n)$ given by

$$(4) \quad \alpha : \Phi(n) \rightarrow \Phi(2n), \quad \alpha(y) = y \quad \text{if } y \text{ odd, else } \alpha(y) = y + n$$

is a group isomorphism. (The proof is a simple exercise, distinguishing 3 cases.) The inverse is

$$\alpha^{-1} : \Phi(2n) \rightarrow \Phi(n), \quad \alpha^{-1}(x) = x \quad \text{if } x < n, \quad \text{else } \alpha^{-1}(x) = x - n.$$

We also need a map β which “halves” the elements of $\Phi(n)$ (modulo n):

$$(5) \quad \beta : \Phi(n) \rightarrow \Phi(n), \quad \beta(y) = y/2 \quad \text{if } y \text{ even, else } \beta(y) = (y + n)/2.$$

The doubling formula for Γ is needed in the following form

$$(7) \quad \Gamma(t) = c_t \Gamma(2t) / \Gamma\left(t + \frac{1}{2}\right),$$

where $c_t = (2\sqrt{\pi})2^{-2t}$. In (7) set $t = x/2n$, where $x \in \Phi(2n)$,

$$(8) \quad \Gamma\left(\frac{x}{2n}\right) = c_{x/2n} \Gamma\left(\frac{x}{n}\right) / \Gamma\left(\frac{x+n}{2n}\right).$$

This equation is of the form

$$(9) \quad \Gamma\left(\frac{x}{2n}\right) = \varepsilon(x) c_{x/2n} \Gamma\left(\frac{y}{n}\right) / \Gamma\left(\frac{z}{n}\right)$$

with $y = x, z = (x + n)/2 \in \Phi(n)$ and $\varepsilon(x) = 1$ when $x < n$. When $x > n$, we apply the reduction formula $\Gamma(t) = (t - 1)\Gamma(t - 1)$ to both Γ 's on the right in (8), and cancel factors $(x - n)/n$. The result is

$$\Gamma\left(\frac{x}{2n}\right) = 2c_{x/2n} \Gamma\left(\frac{x-n}{n}\right) / \Gamma\left(\frac{x-n}{2n}\right),$$

which is of the form (9) with $y = x - n, z = (x - n)/2 \in \Phi(n)$ and $\varepsilon(x) = 2$.

Lemma. Let n be an odd integer, $n > 1$, and $x \in \Phi(2n)$. Then (9) holds, where $y = \alpha^{-1}(x), z = \beta(y) \in \Phi(n)$. Further, $x - n = 2y - 2z$ and $y \equiv 2z \pmod{n}$.

Proof. Distinguish the two cases $x < n$ and $x > n$. (Note that $x \in \Phi(2n)$ is odd.)

If $x < n$, then $\alpha^{-1}(x) = x = y$ (odd), so $\beta(y) = (y + n)/2 = (x + n)/2 = z$.

Also, $2y - 2z = 2x - (x + n) = x - n$ and $2z = y + n \equiv y \pmod{n}$.

If $x > n$, then $\alpha^{-1}(x) = x - n$ (even), so $\beta(y) = y/2 = (x - n)/2 = z$

Also, $2y - 2z = 2(x - n) - (x - n) = x - n$, and $2z = y$.

Proof of (3). The members of $\Phi(n)$ are taken as vertices of a directed labeled graph. The edges are the pairs $(y, z) = (y, \beta(y))$; such an edge is labeled $x = \alpha(y) \in \Phi(2n)$. The vertices have in- and outdegree 1, so the connected components are cycles. In fact, the component of 1 is the cyclic group generated by 2; denote its order by $\nu(n)$. The other components are the cosets. Let B be any one of these cycles. Form the product $P = \prod \Gamma(x/2n)$ of the left sides of (9), where the product extends over $x \in \alpha B$, i.e., the labels of the edges of B . Similarly, take the product of the right sides of (9), and note that the product telescopes as all the Γ 's cancel, leaving only

$$P = \prod_{\alpha B} \varepsilon(x) c_{-x/2n} = (2\sqrt{\pi})^{\nu(n)} 2^{-2(\sum x)/2n} \prod_{\alpha B} \varepsilon(x).$$

Similarly, we have the telescoping sum $\sum(x - n) = \sum(2y - 2z) = 0$, so $\sum x = n\nu(n)$.

Therefore,

$$P = (2\sqrt{\pi})^{\nu(n)} 2^{-2n\nu(n)/2n} \prod_{x \in \alpha B} \varepsilon(x) = \pi^{\nu(n)/2} 2^{b(\alpha B)},$$

where $b(\alpha B)$ is the number of $x \in \alpha B$ that are bigger than n . In summary,

$$\prod_{x \in \alpha B} \Gamma\left(\frac{x}{2n}\right) = 2^{b(\alpha B)} \pi^{\nu(n)/2}.$$

Since α is an isomorphism, $A = \alpha B$ is (a coset of) the subgroup of $\Phi(2n)$ generated by $\alpha(2) = n + 2$. That yields (3).

Corollaries.

1. In Zucker [2] we find, for $n = 2^m - 1$, $m > 1$, that

$$\Gamma\left(\frac{1}{2n}\right) \prod_{k=1}^{m-1} \Gamma\left(\frac{2^k + n}{2n}\right) = 2^{m-1} \pi^{m/2}.$$

Proof: This is a special case of (3). The left side equals

$$\prod_{k=0}^{m-1} \Gamma\left(\frac{\alpha(2^k)}{2n}\right),$$

while $\beta(1) = 2^{m-1}$, $\beta(2^k) = 2^{k-1}$ ($k > 0$). All numerators except for one are $> n$.

2. Complementation. Let A be as in (3), and $A^* = \{2n - x | x \in A\}$, then (3) holds with A^* replacing A , and $b(A^*) = \nu(n) - b(A)$.

Proof: Verify that if $x' \equiv x(n+2) \pmod{2n}$ then $2n - x' \equiv (2n - x)(n+2) \pmod{2n}$.

3. Take the product in (3) over all of $\Phi(2n)$; that is, multiply both sides of (3) as A ranges over the subgroup and its cosets. There are $\varphi(n)/\nu(n)$ choices of A . Each $x \in \Phi(2n)$ greater than n will occur exactly once.

$$\prod_{x \in \Phi(2n)} \Gamma\left(\frac{x}{2n}\right) = (2\pi)^{\varphi(n)/2}.$$

4. Some numerical examples. We determined, for which odd $n < 100$, the number of sets A exceeds 2. There are 9 such values. Among these, only $n = 43$ has 3 A 's, and all of these are self-complementary [2]. In all other cases the number is even, and can be as big as 8. At the other end, there are 16 values of n for which $\nu(n) = \phi(n)$; that can only happen when n is a prime or a prime power.

We list the six sets A for $n = 31$ because only two of them are usually mentioned [2].

$$\begin{aligned} n = 31 : & (1, 33, 35, 39, 47), (3, 17, 37, 43, 55), (5, 9, 41, 49, 51), \\ & (7, 19, 25, 45, 59), (11, 13, 21, 53, 57), (15, 23, 27, 29, 61). \end{aligned}$$

The first one, written out in full, is

$$\Gamma\left(\frac{1}{62}\right) \Gamma\left(\frac{33}{62}\right) \Gamma\left(\frac{35}{62}\right) \Gamma\left(\frac{39}{62}\right) \Gamma\left(\frac{47}{62}\right) = 2^4 \pi^{5/2}.$$

Here, $\nu = 5$, the length of the product, and $b = 4$, the number of numerators bigger than 31. Each numerator, multiplied by 33 (mod 62) yields the next one, in circular order.

In a personal note, H. Chen states the problem of finding minimum sizes of gamma products that have simple values. This paper may be a step in that direction, but any definitive answer will depend on a suitable definition of “simple value”.

References

- [1] Glasser, M. L., Problem 11426, Amer. Math. Monthly, **116**, p. 365, (2009)
- [2] Zucker, L.J., Personal notes (1994)

Department of Mathematics, University of Pennsylvania

Department of Mathematics, University of Washington

nijenhuisalbert@msn.com